

# Fast Quiz #3

## Numerical Functional Analysis

*Præparatus supervivet*

Stefan Engblom

Division of Scientific Computing  
Department of Information Technology  
Uppsala University

Uppsala, May, 2019

## Question 1

**True/False:**  $\|x + y\|^2 \leq \|x\|^2 + \|y\|^2$ .

## Question 2

**True/False:** One way of defining the  $l^p$  inner product is

$$(x, y) = \sum_{j \geq 1} \xi_j^{p/2} \eta_j^{p/2},$$

where  $x = (\xi_j)$  and  $y = (\eta_j)$ .

## Question 3

**True/False:** *Unfortunately,* the norm  $\|x\| = \max_{t \in [a,b]} |x(t)|$  for  $x \in C[a, b]$  can not be obtained from an inner product.

## Question 4

**True/False:** Let  $Y$  be an open subspace of a Hilbert space  $H$ . Then  $Y$  is complete.

## Question 5

**True/False:** Suppose  $(Sv, v) = 0$  for all  $v$  in a complex Hilbert space  $H$ , where  $S$  is a bounded linear operator. Then  $S = 0$ .

## Question 6

**True/False:** A bounded bilinear form  $a(\cdot, \cdot) \geq 0$  defines a norm by  $\|v\|^2 := a(v, v)$ .

## Question 7

**True/False:** If a Hilbert space  $H$  contains a total orthogonal sequence, then  $H$  is separable.



## Question 8

**True/False:** Let  $f \in L^2[0, 1]$ . Assume that  $(e_k)$  is an orthonormal sequence in  $L^2[0, 1]$ . Put

$$\tilde{f} = \sum_{k \geq 1} (f, e_k) e_k.$$

Then  $f = \tilde{f}$ .

## Question 9

**True/False:** If  $x \perp y$ , then  $\|x + y\|^2 = \|x\|^2 + \|y\|^2$ .

## Question 10

**True/False:** Suppose  $(Sv, v) = 0$  for some  $v$  in a complex Hilbert space  $H$ , where  $S$  is a bounded linear operator. Then  $Sv = 0$ .

## Question 11

**True/False:** A bounded coercive bilinear form defines an inner product by  $(u, v) := a(u, v)$ .