



UPPSALA
UNIVERSITET

Linearly constrained Gaussian processes


Niklas Wahlström, Uppsala University, Sweden

Joint work with

Carl Jidling, Philipp Pilar, Thomas Schön, (Uppsala University),
Adrian Wills, Johannes Hendriks, Alexander Gregg, Chris Wensrich (University of Newcastle,
Australia),

SIAM Conference on Uncertainty Quantification,
April 15, 2022


Motivation - Application 1: Magnetic mapping - Indoor localization



Goal: Model magnetic field with a Gaussian process and infer measurements of this field

Question: Can we use any Maxwell's equations to constrain this model?

Motivation - Application 2: Strain field reconstruction



$$y = \mathcal{L}_{\mathbf{x}}\epsilon(\mathbf{x}) + \varepsilon = \frac{1}{L} \int_0^L \hat{\mathbf{n}}^T \epsilon(\mathbf{x}^0 + s\hat{\mathbf{n}}) \hat{\mathbf{n}} ds + \varepsilon$$

Goal: Model $\epsilon(\mathbf{x})$ with a Gaussian process and infer the value of $\epsilon(\mathbf{x}_*)$

Question: Can we use any physical knowledge to constrain this model?

Aim: Introduce constrained Gaussian process regression and demonstrate it on a few examples.

1. GP basics
2. Linear constraints
3. Strain field reconstruction
4. Nonlinear constraints

Distribution over functions

$$\begin{bmatrix} f(\mathbf{x}_1) \\ \vdots \\ f(\mathbf{x}_N) \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mu(\mathbf{x}_1) \\ \vdots \\ \mu(\mathbf{x}_N) \end{bmatrix}, \underbrace{\begin{bmatrix} k(\mathbf{x}_1, \mathbf{x}_1) & \cdots & k(\mathbf{x}_1, \mathbf{x}_N) \\ \vdots & & \vdots \\ k(\mathbf{x}_N, \mathbf{x}_1) & \cdots & k(\mathbf{x}_N, \mathbf{x}_N) \end{bmatrix}}_K \right)$$

Gram matrix

Uniquely specified by mean and covariance function


$$\mu(\mathbf{x}_i) = \mathbb{E}[f(\mathbf{x}_i)]$$

$$k(\mathbf{x}_i, \mathbf{x}_j) = \text{Cov}[f(\mathbf{x}_i), f(\mathbf{x}_j)]$$


Formally

$$f(\mathbf{x}) \sim \mathcal{GP}(\mu(\mathbf{x}), k(\mathbf{x}, \mathbf{x}'))$$


GP basics – linear operator measurements



GP basics – linear operator measurements



GP basics – linear operator measurements



1. GP basics
2. **Linear constraints**
3. Strain field reconstruction
4. Nonlinear constraints

Multivariate GP – constraint incorporation

TOY EXAMPLE

Consider a Gaussian process

$$\mathbf{f}(\mathbf{x}) \sim \mathcal{GP}(\boldsymbol{\mu}(\mathbf{x}), \mathbf{K}(\mathbf{x}, \mathbf{x}'))$$

with two-dimensional input and two-dimensional output

$$\mathbf{f}(\mathbf{x}) = \begin{bmatrix} f_1(\mathbf{x}) \\ f_2(\mathbf{x}) \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$$

Assume that we know from the physics that the all samples from the GP prior should obey the constraint

$$\frac{\partial f_1}{\partial x} + \frac{\partial f_2}{\partial y} = 0 \quad \Leftrightarrow \quad \underbrace{\begin{bmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \end{bmatrix}}_{\mathcal{F}_x} \mathbf{f}(\mathbf{x}) = 0$$

How can we model the covariance function $\mathbf{K}(\mathbf{x}, \mathbf{x}')$ such that this constraint is guaranteed to be obeyed?

Multivariate GP – constraint incorporation

Assume linear constraints

$$\mathcal{F}_x \mathbf{f}(\mathbf{x}) = \mathbf{0}$$

Let $\mathbf{f}(\mathbf{x}) = \mathcal{G}_x \mathbf{g}(\mathbf{x})$, where $\mathbf{g}(\mathbf{x}) \sim \mathcal{GP}(\boldsymbol{\mu}_g(\mathbf{x}), \mathbf{K}_g(\mathbf{x}, \mathbf{x}'))$

$$\mathbf{f}(\mathbf{x}) = \mathcal{G}_x \mathbf{g}(\mathbf{x}) \sim \mathcal{GP}\left(\mathcal{G}_x \boldsymbol{\mu}_g(\mathbf{x}), \mathcal{G}_x \mathbf{K}_g(\mathbf{x}, \mathbf{x}') \mathcal{G}_x^T\right)$$

Then

$$\mathcal{F}_x \mathcal{G}_x \mathbf{g}(\mathbf{x}) = \mathbf{0}$$

Arbitrary $\mathbf{g}(\mathbf{x})$

$$\Rightarrow \mathcal{F}_x \mathcal{G}_x = \mathbf{0}$$

Find \mathcal{G}_x



Carl Jidling, Niklas Wahlström, Adrian Wills, Thomas B. Schön. **Linearly constrained Gaussian processes**. *Advances in Neural Information Processing Systems (NIPS)*, Long Beach, CA, USA, December, 2017.

Multivariate GP – constraint incorporation

TOY EXAMPLE (CONT.)

We consider the function

$$\mathbf{f}(\mathbf{x}) = \begin{bmatrix} f_1(\mathbf{x}) \\ f_2(\mathbf{x}) \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$$

and the constraint

$$\frac{\partial f_1}{\partial x} + \frac{\partial f_2}{\partial y} = 0 \quad \Leftrightarrow \quad \underbrace{\begin{bmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \end{bmatrix}}_{\mathcal{F}_x} \mathbf{f}(\mathbf{x}) = 0$$

Need \mathcal{G}_x such that $\mathcal{F}_x \mathcal{G}_x = \mathbf{0}$. One option is

$$\mathcal{G}_x = \begin{bmatrix} -\frac{\partial}{\partial y} \\ \frac{\partial}{\partial x} \end{bmatrix}$$

since


$$\mathcal{F}_x \mathcal{G}_x = \begin{bmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \end{bmatrix} \begin{bmatrix} -\frac{\partial}{\partial y} \\ \frac{\partial}{\partial x} \end{bmatrix} = -\frac{\partial^2}{\partial x \partial y} + \frac{\partial^2}{\partial y \partial x} = 0.$$

Simulation experiment - toy example

If we choose $k_g(\mathbf{x}, \mathbf{x}') = \sigma_f^2 e^{-\frac{\|\mathbf{x}-\mathbf{x}'\|^2}{2l^2}}$ we get

$$\begin{aligned} \mathbf{K}(\mathbf{x}, \mathbf{x}') &= \mathcal{G}_x \mathcal{G}_{x'}^T k_g(\mathbf{x}, \mathbf{x}') = \begin{bmatrix} -\frac{\partial}{\partial y} \\ \frac{\partial}{\partial x} \end{bmatrix} \begin{bmatrix} -\frac{\partial}{\partial y} & \frac{\partial}{\partial x} \end{bmatrix} k_g(\mathbf{x}, \mathbf{x}') \\ &= \sigma_f^2 e^{-\frac{\|\mathbf{x}-\mathbf{x}'\|^2}{2l^2}} \left(\left(\frac{\mathbf{x}-\mathbf{x}'}{l} \right) \left(\frac{\mathbf{x}-\mathbf{x}'}{l} \right)^T - \left(1 - \frac{\|\mathbf{x}-\mathbf{x}'\|^2}{l^2} \right) I_2 \right) \end{aligned}$$



Below we have simulated a field which we know fulfills the constraint



1. GP basics
2. Linear constraints
3. Strain field reconstruction
4. Nonlinear constraints

Strain field reconstruction


Deformed object



Reconstruct the *strain tensor*

$$\epsilon(\mathbf{x}) = \begin{bmatrix} \epsilon_{xx}(\mathbf{x}) & \epsilon_{xy}(\mathbf{x}) \\ \epsilon_{xy}(\mathbf{x}) & \epsilon_{yy}(\mathbf{x}) \end{bmatrix}$$


Strain field reconstruction - Bragg-edge analysis



$$y = \frac{1}{L} \int_0^L \hat{\mathbf{n}}^T \boldsymbol{\epsilon}(\mathbf{x}^0 + s\hat{\mathbf{n}}) \hat{\mathbf{n}} ds + \varepsilon$$

$$\hat{\mathbf{n}} = \begin{bmatrix} n_x \\ n_y \end{bmatrix}, \quad \varepsilon \sim \mathcal{N}(0, \sigma^2)$$

Strain field reconstruction - Bragg-edge analysis



Vectorised form

$$y = \frac{1}{L} \int_0^L \vec{\mathbf{n}}^T \mathbf{f}(\mathbf{x}^0 + s\hat{\mathbf{n}}) ds + \varepsilon = \mathcal{L}_{\mathbf{x}}[\mathbf{f}(\mathbf{x})] + \varepsilon$$

$$\mathbf{f}(\mathbf{x}) = \begin{bmatrix} f_{xx}(\mathbf{x}) \\ f_{xy}(\mathbf{x}) \\ f_{yy}(\mathbf{x}) \end{bmatrix} = \begin{bmatrix} \epsilon_{xx}(\mathbf{x}) \\ \epsilon_{xy}(\mathbf{x}) \\ \epsilon_{yy}(\mathbf{x}) \end{bmatrix}, \quad \vec{\mathbf{n}} = \begin{bmatrix} n_x^2 \\ 2n_x n_y \\ n_y^2 \end{bmatrix}$$

Strain field reconstruction – constraint incorporation

A physical strain field must satisfy the *equilibrium constraints*

$$0 = \frac{\partial f_{xx}(\mathbf{x})}{\partial x} + (1 - \nu) \frac{\partial f_{xy}(\mathbf{x})}{\partial y} + \nu \frac{\partial f_{yy}(\mathbf{x})}{\partial x}$$
$$0 = \nu \frac{\partial f_{xx}(\mathbf{x})}{\partial y} + (1 - \nu) \frac{\partial f_{xy}(\mathbf{x})}{\partial x} + \frac{\partial f_{yy}(\mathbf{x})}{\partial y}$$

These can be written as

$$\mathbf{0} = \underbrace{\begin{bmatrix} \frac{\partial}{\partial x} & (1 - \nu) \frac{\partial}{\partial y} & \nu \frac{\partial}{\partial x} \\ \nu \frac{\partial}{\partial y} & (1 - \nu) \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \end{bmatrix}}_{\mathcal{F}_x} \mathbf{f}(\mathbf{x})$$

Strain field reconstruction – constraint incorporation

We get

$$\mathcal{G}_{\mathbf{x}} = \begin{bmatrix} \frac{\partial^2}{\partial y^2} - \nu \frac{\partial^2}{\partial x^2} \\ -(1 + \nu) \frac{\partial^2}{\partial x \partial y} \\ \frac{\partial^2}{\partial x^2} - \nu \frac{\partial^2}{\partial y^2} \end{bmatrix}$$

Hence

$$\mathbf{f}(\mathbf{x}) = \mathcal{G}_{\mathbf{x}} \varphi(\mathbf{x})$$

Now let

$$\varphi(\mathbf{x}) \sim \mathcal{GP}(0, k_{\varphi}(\mathbf{x}, \mathbf{x}'))$$


Then

$$\mathbf{f}(\mathbf{x}) \sim \mathcal{GP}\left(0, \mathcal{G}_{\mathbf{x}} \mathcal{G}_{\mathbf{x}'}^{\top} k_{\varphi}(\mathbf{x}, \mathbf{x}')\right)$$

Note

$$y = \mathcal{L}_{\mathbf{x}}[\mathcal{G}_{\mathbf{x}} \varphi(\mathbf{x})] + \varepsilon$$

Strain field reconstruction – experimental results



1. GP basics
2. Linear constraints
3. Strain field reconstruction
4. Nonlinear constraints

Nonlinearly constrained Gaussian processes- idea

Question: What can we do if we have nonlinear constraints?

We focus on sum-constrained Gaussian processes

$$\mathcal{F}[\mathbf{f}(\mathbf{x})] = \sum_i a_i h_i(f_i(\mathbf{x})) = C,$$

where $h_i(\cdot)$ is a non-linear function.

Idea: Transform outputs $f'_i = h_i(f_i)$. The constraint will then be linear

$$\mathcal{F}[\mathbf{f}'(\mathbf{x})] = \sum a_i f'_i(\mathbf{x}) = C,$$

Let f'_i be the output of the GP and train it on transform data $y'_i = h_i(y_i)$.

Nonlinearly constrained Gaussian processes- toy example

TOY EXAMPLE (HARMONIC OSCILLATOR)

Motion modelled by multitask Gaussian process

$$\mathbf{f}(t) = \begin{bmatrix} z(t) \\ v(t) \end{bmatrix} \quad \begin{array}{l} z(t) : \text{displacement} \\ v(t) : \text{velocity} \end{array}$$

Constraint: energy conservation (friction neglected)


$$E = E_{\text{pot}}(t) + E_{\text{kin}}(t) = kz(t)^2/2 + mv(t)^2/2,$$

Sum constraint parameters




$$\mathcal{F}[\mathbf{f}(\mathbf{x})] = \sum_i a_i h_i(f_i(\mathbf{x})) = C,$$

$$a_1 = k/2, \quad h_1(z) = z^2$$

$$a_2 = m/2, \quad h_2(v) = v^2, \quad C = E.$$



Nonlinearly constrained Gaussian processes - toy example



Left: Results for unconstrained GP

Middle: Results for transformed output learned by the constrained GP

Right: The back transformed output. The results for the unconstrained GP are used to recover the signs.

Nonlinearly constrained Gaussian processes


- Double pendulum (real data)

REAL DATA EXAMPLE (DOUBLE PENDULUM)

We model both positions z_x, z_y and velocities v_x, v_y of the two masses, (i.e. 8 outputs), while at the same time respecting the law of energy conservation


$$E = m_b g z_{by} + m_g g z_{gy} + \frac{m_b}{2} (v_{bx}^2 + v_{by}^2) + \frac{m_g}{2} (v_{gx}^2 + v_{gy}^2),$$

Indices b and g refer to blue and green pendulum, respectively.



Nonlinearly constrained Gaussian processes

- Double pendulum (real data) - Results



Left: Results for unconstrained GP **Right:** Results for constrained GP



Conclusions and references

- ▶ Linear constraints can be incorporated in Gaussian processes
- ▶ Promising results on simulated and real data experiments
- ▶ The idea can also be extended to a nonlinear constraints

References



Carl Jidling, Niklas Wahlstöm, Adrian Wills, Thomas B. Schön. **Linearly constrained Gaussian processes**. *Advances in Neural Information Processing Systems (NIPS)*, Long Beach, CA, USA, December, 2017.



Arno Solin, Manon Kok, Niklas Wahlström, Thomas B. Schön, and Simo Särkkä. **Modeling and interpolation of the ambient magnetic field by Gaussian processes**. *IEEE Transactions on Robotics*, 34(4):1112 – 1127, 2018



Carl Jidling, Johannes Hendriks, Niklas Wahlström, Alexander Gregg, Thomas B. Schön, Chris Wensrich, Adrian Wills. **Probabilistic modelling and reconstruction of strain**, *Nuclear instruments and methods in physics research section B*, 436:141-155, 2018.



Philipp Pilar, Carl Jidling, Thomas B. Schön, Niklas Wahlström. **Incorporating sum constraints into multitask Gaussian processes**, *arXiv preprint arXiv:1504.00702*, 2022.