



UPPSALA
UNIVERSITET

Constrained Gaussian processes for strain field reconstruction

Niklas Wahlström, Uppsala University

Joint work with

Carl Jidling, Thomas Schön, (Uppsala University),

Adrian Wills, Johannes Hendriks, Alexander Gregg, Chris Wensrich (University of Newcastle, Australia),


Physics-informed machine learning – The 3rd Sheffield Workshop on Structural Dynamics
December 8, 2020.

Tomography intuition

				11	22
			↗ ↘		13
5	9	4	↗ ↘		18
2	7	12	→		21
11	1	4	↘ ↙		16
↓ ↓ ↓ ↘			↘		21
28	17	20	3		16

				3	16
			↗ ↘		16
?	?	?	↗ ↘		19
?	?	?	→		10
?	?	?	↘ ↙		25
↓ ↓ ↓ ↘			↘		8
21	15	18	11		22

Application: Strain field reconstruction



$$y = \mathcal{L}_{\mathbf{x}}\epsilon(\mathbf{x}) + \varepsilon = \frac{1}{L} \int_0^L \hat{\mathbf{n}}^\top \epsilon(\mathbf{x}^0 + s\hat{\mathbf{n}}) \hat{\mathbf{n}} ds + \varepsilon$$

Goal: Model $\epsilon(\mathbf{x})$ with a Gaussian process and infer the value of $\epsilon(\mathbf{x}_*)$

Question: Can we use any physical knowledge to constrain this model?

Aim: Introduce Gaussian process regression and demonstrate its potential for tomographic reconstruction

1. Intuition
2. GP basics
3. Linear constraints
4. Strain revisited

				3	16
					16
					19
					10
					25
					8
21	15	18	11	22	

Distribution over functions

$$\begin{bmatrix} f(\mathbf{x}_1) \\ \vdots \\ f(\mathbf{x}_N) \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mu(\mathbf{x}_1) \\ \vdots \\ \mu(\mathbf{x}_N) \end{bmatrix}, \underbrace{\begin{bmatrix} k(\mathbf{x}_1, \mathbf{x}_1) & \cdots & k(\mathbf{x}_1, \mathbf{x}_N) \\ \vdots & & \vdots \\ k(\mathbf{x}_N, \mathbf{x}_1) & \cdots & k(\mathbf{x}_N, \mathbf{x}_N) \end{bmatrix}}_K \right)$$

Gram matrix

Uniquely specified by mean and covariance function

$$\mu(\mathbf{x}_i) = \mathbb{E}[f(\mathbf{x}_i)]$$

$$k(\mathbf{x}_i, \mathbf{x}_j) = \text{Cov}[f(\mathbf{x}_i), f(\mathbf{x}_j)]$$

Formally

$$f(\mathbf{x}) \sim \mathcal{GP}(\mu(\mathbf{x}), k(\mathbf{x}, \mathbf{x}'))$$

GP basics – prediction

Let

$$\mathbf{y} = [y_1, y_2 \dots, y_N]^T$$

Then

$$\begin{bmatrix} \mathbf{y} \\ f(\mathbf{x}_*) \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{0} \\ 0 \end{bmatrix}, \begin{bmatrix} K + \sigma^2 I & \mathbf{k} \\ \mathbf{k}^T & k(\mathbf{x}_*, \mathbf{x}_*) \end{bmatrix} \right)$$

$$K_{ij} = k(\mathbf{x}_i, \mathbf{x}_j)$$

$$\mathbf{k}_i = k(\mathbf{x}_i, \mathbf{x}_*)$$

and

$$\mathbb{E}[f(\mathbf{x}_*)|\mathbf{y}] = \mathbf{k}^T (K + \sigma^2 I)^{-1} \mathbf{y}$$

$$\mathbb{V}[f(\mathbf{x}_*)|\mathbf{y}] = k(\mathbf{x}_*, \mathbf{x}_*) - \mathbf{k}^T (K + \sigma^2 I)^{-1} \mathbf{k}$$

GP basics – linear operator measurements

Linear operator measurements

$$y = \mathcal{L}_{\mathbf{x}} f(\mathbf{x}) + \varepsilon$$

Then

$$\mathbb{E}[f(\mathbf{x}_*) | \mathbf{y}] = \mathbf{q}^\top (Q + \sigma^2 I)^{-1} \mathbf{y}$$

$$\mathbb{V}[f(\mathbf{x}_*) | \mathbf{y}] = k(\mathbf{x}_*, \mathbf{x}_*) - \mathbf{q}^\top (Q + \sigma^2 I)^{-1} \mathbf{q}$$

where


$$Q_{ij} = \mathcal{L}_{\mathbf{x}_i} \mathcal{L}_{\mathbf{x}_j} k(\mathbf{x}_i, \mathbf{x}_j)$$

$$\mathbf{q}_i = \mathcal{L}_{\mathbf{x}_i} k(\mathbf{x}_i, \mathbf{x}_*)$$


Example:

$$y_i = \int_{a_i}^{b_i} f(x) dx \Rightarrow \begin{cases} Q_{ij} = \int_{a_i}^{b_i} \int_{a_j}^{b_j} k(x, x') dx' dx \\ \mathbf{q}_i = \int_{a_i}^{b_i} k(x, x_*) dx \end{cases}$$

GP basics – linear operator measurements



GP basics – linear operator measurements



Outline

1. Intuition
2. GP basics
- 3. Linear constraints**
4. Strain revisited

				3	16
					16
					19
					10
					25
					8
21	15	18	11	22	

Multivariate GP – constraint incorporation

TOY EXAMPLE

Consider a Gaussian process

$$f(\mathbf{x}) \sim \mathcal{GP}(\boldsymbol{\mu}(\mathbf{x}), K(\mathbf{x}, \mathbf{x}'))$$

with two-dimensional input and two-dimensional output

$$\mathbf{f}(\mathbf{x}) = \begin{bmatrix} f_1(\mathbf{x}) \\ f_2(\mathbf{x}) \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$$

Assume that we know from the physics that all samples from the GP prior should obey the constraint

$$\frac{\partial f_1}{\partial x} + \frac{\partial f_2}{\partial y} = 0 \quad \Leftrightarrow \quad \underbrace{\begin{bmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \end{bmatrix}}_{\mathcal{F}_x} \mathbf{f}(\mathbf{x}) = 0$$

How can we model the covariance function $\mathbf{K}(\mathbf{x}, \mathbf{x}')$ such that this constraint is guaranteed to be obeyed?

Multivariate GP – constraint incorporation

Assume linear constraints

$$\mathcal{F}_x f(\mathbf{x}) = \mathbf{0}$$

Let $f(\mathbf{x}) = \mathcal{G}_x g(\mathbf{x})$

$$f(\mathbf{x}) = \mathcal{G}_x g(\mathbf{x}) \sim \mathcal{GP} \left(\mathcal{G}_x \mu_g(\mathbf{x}), \mathcal{G}_x \mathbf{K}_g(\mathbf{x}, \mathbf{x}') \mathcal{G}_x^T \right)$$

Then

$$\mathcal{F}_x \mathcal{G}_x g(\mathbf{x}) = \mathbf{0}$$

Arbitrary $g(\mathbf{x})$

$$\Rightarrow \mathcal{F}_x \mathcal{G}_x = \mathbf{0}$$

Find \mathcal{G}_x



Carl Jidling, Niklas Wahlström, Adrian Wills, Thomas B. Schön. **Linearly constrained Gaussian processes**. *Advances in Neural Information Processing Systems (NIPS)*, Long Beach, CA, USA, December, 2017.

Multivariate GP – constraint incorporation

TOY EXAMPLE (CONT.)

We consider the function

$$\mathbf{f}(\mathbf{x}) = \begin{bmatrix} f_1(\mathbf{x}) \\ f_2(\mathbf{x}) \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$$

and the constraint

$$\frac{\partial f_1}{\partial x} + \frac{\partial f_2}{\partial y} = 0 \quad \Leftrightarrow \quad \underbrace{\begin{bmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \end{bmatrix}}_{\mathcal{F}_x} \mathbf{f}(\mathbf{x}) = 0$$

Need \mathcal{G}_x such that $\mathcal{F}_x \mathcal{G}_x = \mathbf{0}$. One option is

$$\mathcal{G}_x = \begin{bmatrix} -\frac{\partial}{\partial y} \\ \frac{\partial}{\partial x} \end{bmatrix}$$

since

$$\mathcal{F}_x \mathcal{G}_x = \begin{bmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \end{bmatrix} \begin{bmatrix} -\frac{\partial}{\partial y} \\ \frac{\partial}{\partial x} \end{bmatrix} = -\frac{\partial^2}{\partial x \partial y} + \frac{\partial^2}{\partial y \partial x} = 0.$$

Algorithm idea – toy example

Step 1: Assume that \mathcal{G}_x contains the same operators as \mathcal{F}_x

$$\mathcal{G}_x = \begin{bmatrix} \gamma_{11} & \gamma_{12} \\ \gamma_{21} & \gamma_{22} \end{bmatrix} \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{bmatrix}$$

Step 2: Expand

$$\begin{aligned} \mathcal{F}_x \mathcal{G}_x &= \begin{bmatrix} \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \end{bmatrix} \begin{bmatrix} \gamma_{11} & \gamma_{12} \\ \gamma_{21} & \gamma_{22} \end{bmatrix} \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{bmatrix} \\ &= \gamma_{11} \frac{\partial^2}{\partial x^2} + (\gamma_{12} + \gamma_{21}) \frac{\partial^2}{\partial x \partial y} + \gamma_{22} \frac{\partial^2}{\partial y^2} \end{aligned}$$

Algorithm idea – toy example

Step 3: We need

$$\begin{cases} \gamma_{11} = 0 \\ \gamma_{12} = -\gamma_{21} \\ \gamma_{22} = 0 \end{cases}$$

Step 4: Choosing $\gamma_{21} = 1$, we get

$$\mathcal{G}_x = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \frac{\partial}{\partial x} \\ \frac{\partial}{\partial y} \end{bmatrix} = \begin{bmatrix} -\frac{\partial}{\partial y} \\ \frac{\partial}{\partial x} \end{bmatrix}$$

No solution? Retry with higher order operators!

Even more formal treatment based on polynomial rings and Gröbner basis theory is published in




Simulation experiment - toy example

If we choose $k_g(\mathbf{x}, \mathbf{x}') = \sigma_f^2 e^{-\frac{\|\mathbf{x}-\mathbf{x}'\|^2}{2l^2}}$ we get

$$\begin{aligned}\mathbf{K}(\mathbf{x}, \mathbf{x}') &= \mathcal{G}_x \mathcal{G}_{x'}^T k_g(\mathbf{x}, \mathbf{x}') = \begin{bmatrix} -\frac{\partial}{\partial y} \\ \frac{\partial}{\partial x} \end{bmatrix} \begin{bmatrix} -\frac{\partial}{\partial y} & \frac{\partial}{\partial x} \end{bmatrix} k_g(\mathbf{x}, \mathbf{x}') \\ &= \sigma_f^2 e^{-\frac{\|\mathbf{x}-\mathbf{x}'\|^2}{2l^2}} \left(\left(\frac{\mathbf{x}-\mathbf{x}'}{l} \right) \left(\frac{\mathbf{x}-\mathbf{x}'}{l} \right)^T - \left(1 - \frac{\|\mathbf{x}-\mathbf{x}'\|^2}{l^2} \right) I_2 \right)\end{aligned}$$

Below we have simulated a field which we know fulfills the constraint





Outline

1. Intuition
2. GP basics
3. Linear constraints
4. Strain revisited

				3	16
					16
					19
					10
					25
					8
21	15	18	11	22	

Strain field reconstruction

Deformed object



Reconstruct the *strain tensor*

$$\epsilon(\mathbf{x}) = \begin{bmatrix} \epsilon_{xx}(\mathbf{x}) & \epsilon_{xy}(\mathbf{x}) \\ \epsilon_{xy}(\mathbf{x}) & \epsilon_{yy}(\mathbf{x}) \end{bmatrix}$$




Strain field reconstruction

Bragg's law


$$\lambda = 2d \sin \theta$$

Average strain

$$\langle \epsilon \rangle = \frac{d - d_0}{d_0}$$




Strain field reconstruction



$$y = \frac{1}{L} \int_0^L \hat{\mathbf{n}}^T \boldsymbol{\epsilon}(\mathbf{x}^0 + s\hat{\mathbf{n}}) \hat{\mathbf{n}} ds + \varepsilon$$

$$\hat{\mathbf{n}} = \begin{bmatrix} n_x \\ n_y \end{bmatrix}, \quad \varepsilon \sim \mathcal{N}(0, \sigma^2)$$

Strain field reconstruction



Vectorised form

$$y = \frac{1}{L} \int_0^L \vec{\mathbf{n}}^T \mathbf{f}(\mathbf{x}^0 + s\hat{\mathbf{n}}) ds + \varepsilon$$

$$\mathbf{f}(\mathbf{x}) = \begin{bmatrix} f_{xx}(\mathbf{x}) \\ f_{xy}(\mathbf{x}) \\ f_{yy}(\mathbf{x}) \end{bmatrix} = \begin{bmatrix} \epsilon_{xx}(\mathbf{x}) \\ \epsilon_{xy}(\mathbf{x}) \\ \epsilon_{yy}(\mathbf{x}) \end{bmatrix}, \quad \vec{\mathbf{n}} = \begin{bmatrix} n_x^2 \\ 2n_x n_y \\ n_y^2 \end{bmatrix}$$

Strain field reconstruction – prediction

Put a GP on $\mathbf{f}(\mathbf{x})$

$$\mathbf{f}(\mathbf{x}) \sim \mathcal{GP}(\mathbf{0}, \mathbf{K}(\mathbf{x}, \mathbf{x}'))$$

As before

$$\mathbb{E}[\mathbf{f}(\mathbf{x}_*)|\mathbf{y}] = \mathbf{Q}_*^T (\mathbf{Q} + \sigma^2 \mathbf{I})^{-1} \mathbf{y}$$

$$\text{Cov}[\mathbf{f}(\mathbf{x}_*)|\mathbf{y}] = \mathbf{K}(\mathbf{x}_*, \mathbf{x}_*) - \mathbf{Q}_*^T (\mathbf{Q} + \sigma^2 \mathbf{I})^{-1} \mathbf{Q}_*$$

$$Q_{ij} = \frac{1}{L_i L_j} \int_0^{L_i} \int_0^{L_j} \vec{\mathbf{n}}_i^T \mathbf{K}(\mathbf{x}_i^0 + s \hat{\mathbf{n}}_i, \mathbf{x}_j^0 + t \hat{\mathbf{n}}_j) \vec{\mathbf{n}}_j dt ds$$

$$(Q_*)_i = \frac{1}{L_i} \int_0^{L_i} \vec{\mathbf{n}}_i^T \mathbf{K}(\mathbf{x}_i^0 + s \hat{\mathbf{n}}_i, \mathbf{x}_*) ds$$

Strain field reconstruction – covariance model

Since $\mathbf{f}(\mathbf{x})$ is multivariate, the covariance function is a *matrix*

$$\mathbf{K}(\mathbf{x}, \mathbf{x}') = \begin{bmatrix} k_{11}(\mathbf{x}, \mathbf{x}') & k_{12}(\mathbf{x}, \mathbf{x}') & k_{13}(\mathbf{x}, \mathbf{x}') \\ k_{21}(\mathbf{x}, \mathbf{x}') & k_{22}(\mathbf{x}, \mathbf{x}') & k_{23}(\mathbf{x}, \mathbf{x}') \\ k_{31}(\mathbf{x}, \mathbf{x}') & k_{32}(\mathbf{x}, \mathbf{x}') & k_{33}(\mathbf{x}, \mathbf{x}') \end{bmatrix}$$

How should we select $\mathbf{K}(\mathbf{x}, \mathbf{x}')$?

Strain field reconstruction – constraint incorporation

A physical strain field must satisfy the *equilibrium constraints*

$$0 = \frac{\partial f_{xx}(\mathbf{x})}{\partial x} + (1 - \nu) \frac{\partial f_{xy}(\mathbf{x})}{\partial y} + \nu \frac{\partial f_{yy}(\mathbf{x})}{\partial x}$$
$$0 = \nu \frac{\partial f_{xx}(\mathbf{x})}{\partial y} + (1 - \nu) \frac{\partial f_{xy}(\mathbf{x})}{\partial x} + \frac{\partial f_{yy}(\mathbf{x})}{\partial y}$$

These can be written as

$$\mathbf{0} = \underbrace{\begin{bmatrix} \frac{\partial}{\partial x} & (1 - \nu) \frac{\partial}{\partial y} & \nu \frac{\partial}{\partial x} \\ \nu \frac{\partial}{\partial y} & (1 - \nu) \frac{\partial}{\partial x} & \frac{\partial}{\partial y} \end{bmatrix}}_{\mathcal{F}_x} \mathbf{f}(\mathbf{x}) = \begin{bmatrix} \mathbf{c}_1^\top \\ \mathbf{c}_2^\top \end{bmatrix} \mathbf{f}(\mathbf{x})$$

Strain field reconstruction – constraint incorporation

We get

$$\mathcal{G}_{\mathbf{x}} = \mathbf{c}_1 \times \mathbf{c}_2 = \begin{bmatrix} \frac{\partial^2}{\partial y^2} - \nu \frac{\partial^2}{\partial x^2} \\ -(1 + \nu) \frac{\partial^2}{\partial x \partial y} \\ \frac{\partial^2}{\partial x^2} - \nu \frac{\partial^2}{\partial y^2} \end{bmatrix}$$

Hence

$$\mathbf{f}(\mathbf{x}) = \mathcal{G}_{\mathbf{x}} \varphi(\mathbf{x})$$

The scalar $\varphi(\mathbf{x})$ is the *Airy stress function*. Now let

$$\varphi(\mathbf{x}) \sim \mathcal{GP}(0, k_{\varphi}(\mathbf{x}, \mathbf{x}'))$$


Then

$$\mathbf{f}(\mathbf{x}) \sim \mathcal{GP}\left(0, \mathcal{G}_{\mathbf{x}} \mathcal{G}_{\mathbf{x}'}^{\top} k_{\varphi}(\mathbf{x}, \mathbf{x}')\right)$$

Note

$$y = \mathcal{L}_{\mathbf{x}}[\mathcal{G}_{\mathbf{x}} \varphi(\mathbf{x})] + \varepsilon$$

Strain field reconstruction – experimental results



The Gaussian process is well suited for tomographic reconstruction

- ▶ Linear transformations are easily incorporated
- ▶ Physical laws can be built into the model
- ▶ Promising results on real data experiments

The combination of model driven **physical knowledge** and data driven **flexibility** is promising